

## In vivo demonstration that $\alpha$ synuclein oligomers are toxic.

**Journal:** Proc Natl Acad Sci U S A

**Publication Year:** 2011

**Authors:** B Winner, R Jappelli, S K Maji, P A Desplats, L Boyer, S Aigner, C Hetzer, T Loher, M Vilar, S Campioni, C Tzitzilonis, A Soragni, S Jessberger, H Mira, A Consiglio, E Pham, E Masliah, F H Gage, R Riek

**PubMed link:** 21325059

**Funding Grants:** Interdisciplinary Stem Cell Training Program at UCSD II

### Public Summary:

The aggregation of proteins into oligomers and amyloid fibrils is characteristic of several neurodegenerative diseases, including Parkinson disease (PD). In PD, the process of aggregation of  $\alpha$ -synuclein ( $\alpha$ -syn) from monomers, via oligomeric intermediates, into amyloid fibrils is considered the disease-causative toxic mechanism. We developed  $\alpha$ -syn mutants that promote oligomer or fibril formation and tested the toxicity of these mutants by using a rat lentivirus system to investigate loss of dopaminergic neurons in the substantia nigra. The most severe dopaminergic loss in the substantia nigra is observed in animals with the  $\alpha$ -syn variants that form oligomers (i.e., E57K and E35K), whereas the  $\alpha$ -syn variants that form fibrils very quickly are less toxic. We show that  $\alpha$ -syn oligomers are toxic in vivo and that  $\alpha$ -syn oligomers might interact with and potentially disrupt membranes.

### Scientific Abstract:

The aggregation of proteins into oligomers and amyloid fibrils is characteristic of several neurodegenerative diseases, including Parkinson disease (PD). In PD, the process of aggregation of alpha-synuclein (alpha-syn) from monomers, via oligomeric intermediates, into amyloid fibrils is considered the disease-causative toxic mechanism. We developed alpha-syn mutants that promote oligomer or fibril formation and tested the toxicity of these mutants by using a rat lentivirus system to investigate loss of dopaminergic neurons in the substantia nigra. The most severe dopaminergic loss in the substantia nigra is observed in animals with the alpha-syn variants that form oligomers (i.e., E57K and E35K), whereas the alpha-syn variants that form fibrils very quickly are less toxic. We show that alpha-syn oligomers are toxic in vivo and that alpha-syn oligomers might interact with and potentially disrupt membranes.

### PNAS Lens Free Article Link:



**Source URL:** <https://www.cirm.ca.gov/about-cirm/publications/vivo-demonstration-alpha-synuclein-oligomers-are-toxic>